
Desk Reference
to the
Diagnostic Criteria
From

DSM-5™
(2013)



American
Psychiatric
Publishing
A Division of American Psychiatric Association

Washington, DC
London, England

6. **Koro:** Related to *dhat syndrome* (see "Glossary of Cultural Concepts of Distress" in the Appendix to DSM-5), an episode of sudden and intense anxiety that the penis (or the vulva and nipples in females) will recede into the body, possibly leading to death.
7. **Jikoshu-kyofu:** A variant of *taijin kyofusho* (see "Glossary of Cultural Concepts of Distress" in the Appendix to DSM-5) characterized by fear of having an offensive body odor (also termed *olfactory reference syndrome*).

Unspecified Obsessive-Compulsive and Related Disorder

300.3 (F42)

This category applies to presentations in which symptoms characteristic of an obsessive-compulsive and related disorder that cause clinically significant distress or impairment in social, occupational, or other important areas of functioning predominate but do not meet the full criteria for any of the disorders in the obsessive-compulsive and related disorders diagnostic class. The unspecified obsessive-compulsive and related disorder category is used in situations in which the clinician chooses *not* to specify the reason that the criteria are not met for a specific obsessive-compulsive and related disorder, and includes presentations in which there is insufficient information to make a more specific diagnosis (e.g., in emergency room settings).

Trauma- and Stressor-Related Disorders

Reactive Attachment Disorder

313.89 (F94.1)

- A. A consistent pattern of inhibited, emotionally withdrawn behavior toward adult caregivers, manifested by both of the following:
1. The child rarely or minimally seeks comfort when distressed.
 2. The child rarely or minimally responds to comfort when distressed.
- B. A persistent social and emotional disturbance characterized by at least two of the following:
1. Minimal social and emotional responsiveness to others.
 2. Limited positive affect.
 3. Episodes of unexplained irritability, sadness, or fearfulness that are evident even during nonthreatening interactions with adult caregivers.
- C. The child has experienced a pattern of extremes of insufficient care as evidenced by at least one of the following:
1. Social neglect or deprivation in the form of persistent lack of having basic emotional needs for comfort, stimulation, and affection met by caregiving adults.
 2. Repeated changes of primary caregivers that limit opportunities to form stable attachments (e.g., frequent changes in foster care).
 3. Rearing in unusual settings that severely limit opportunities to form selective attachments (e.g., institutions with high child-to-caregiver ratios).
- D. The care in Criterion C is presumed to be responsible for the disturbed behavior in Criterion A (e.g., the disturbances in Criterion A began following the lack of adequate care in Criterion C).

- E. The criteria are not met for autism spectrum disorder.
- F. The disturbance is evident before age 5 years.
- G. The child has a developmental age of at least 9 months.

Specify if:

Persistent: The disorder has been present for more than 12 months.

Specify current severity:

Reactive attachment disorder is specified as **severe** when a child exhibits all symptoms of the disorder, with each symptom manifesting at relatively high levels.

Disinhibited Social Engagement Disorder

313.89 (F94.2)

- A. A pattern of behavior in which a child actively approaches and interacts with unfamiliar adults and exhibits at least two of the following:
 1. Reduced or absent reticence in approaching and interacting with unfamiliar adults.
 2. Overly familiar verbal or physical behavior (that is not consistent with culturally sanctioned and with age-appropriate social boundaries).
 3. Diminished or absent checking back with adult caregiver after venturing away, even in unfamiliar settings.
 4. Willingness to go off with an unfamiliar adult with minimal or no hesitation.
- B. The behaviors in Criterion A are not limited to impulsivity (as in attention-deficit/hyperactivity disorder) but include socially disinhibited behavior.
- C. The child has experienced a pattern of extremes of insufficient care as evidenced by at least one of the following:
 1. Social neglect or deprivation in the form of persistent lack of having basic emotional needs for comfort, stimulation, and affection met by caregiving adults.
 2. Repeated changes of primary caregivers that limit opportunities to form stable attachments (e.g., frequent changes in foster care).

- 3. Rearing in unusual settings that severely limit opportunities to form selective attachments (e.g., institutions with high child-to-caregiver ratios).
- D. The care in Criterion C is presumed to be responsible for the disturbed behavior in Criterion A (e.g., the disturbances in Criterion A began following the pathogenic care in Criterion C).
- E. The child has a developmental age of at least 9 months.

Specify if:

Persistent: The disorder has been present for more than 12 months.

Specify current severity:

Disinhibited social engagement disorder is specified as **severe** when the child exhibits all symptoms of the disorder, with each symptom manifesting at relatively high levels.

Posttraumatic Stress Disorder

309.81 (F43.10)

Posttraumatic Stress Disorder

Note: The following criteria apply to adults, adolescents, and children older than 6 years. For children 6 years and younger, see corresponding criteria below.

- A. Exposure to actual or threatened death, serious injury, or sexual violence in one (or more) of the following ways:
 1. Directly experiencing the traumatic event(s).
 2. Witnessing, in person, the event(s) as it occurred to others.
 3. Learning that the traumatic event(s) occurred to a close family member or close friend. In cases of actual or threatened death of a family member or friend, the event(s) must have been violent or accidental.
 4. Experiencing repeated or extreme exposure to aversive details of the traumatic event(s) (e.g., first responders collecting human remains; police officers repeatedly exposed to details of child abuse).

Note: Criterion A4 does not apply to exposure through electronic media, television, movies, or pictures, unless this exposure is work related.

- B. Presence of one (or more) of the following intrusion symptoms associated with the traumatic event(s), beginning after the traumatic event(s) occurred:
1. Recurrent, involuntary, and intrusive distressing memories of the traumatic event(s).
Note: In children older than 6 years, repetitive play may occur in which themes or aspects of the traumatic event(s) are expressed.
 2. Recurrent distressing dreams in which the content and/or affect of the dream are related to the traumatic event(s).
Note: In children, there may be frightening dreams without recognizable content.
 3. Dissociative reactions (e.g., flashbacks) in which the individual feels or acts as if the traumatic event(s) were recurring. (Such reactions may occur on a continuum, with the most extreme expression being a complete loss of awareness of present surroundings.)
Note: In children, trauma-specific reenactment may occur in play.
 4. Intense or prolonged psychological distress at exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).
 5. Marked physiological reactions to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).
- C. Persistent avoidance of stimuli associated with the traumatic event(s), beginning after the traumatic event(s) occurred, as evidenced by one or both of the following:
1. Avoidance of or efforts to avoid distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).
 2. Avoidance of or efforts to avoid external reminders (people, places, conversations, activities, objects, situations) that

arouse distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).

- D. Negative alterations in cognitions and mood associated with the traumatic event(s), beginning or worsening after the traumatic event(s) occurred, as evidenced by two (or more) of the following:
1. Inability to remember an important aspect of the traumatic event(s) (typically due to dissociative amnesia and not to other factors such as head injury, alcohol, or drugs).
 2. Persistent and exaggerated negative beliefs or expectations about oneself, others, or the world (e.g., "I am bad," "No one can be trusted," "The world is completely dangerous," "My whole nervous system is permanently ruined").
 3. Persistent, distorted cognitions about the cause or consequences of the traumatic event(s) that lead the individual to blame himself/herself or others.
 4. Persistent negative emotional state (e.g., fear, horror, anger, guilt, or shame).
 5. Markedly diminished interest or participation in significant activities.
 6. Feelings of detachment or estrangement from others.
 7. Persistent inability to experience positive emotions (e.g., inability to experience happiness, satisfaction, or loving feelings).
- E. Marked alterations in arousal and reactivity associated with the traumatic event(s), beginning or worsening after the traumatic event(s) occurred, as evidenced by two (or more) of the following:
1. Irritable behavior and angry outbursts (with little or no provocation) typically expressed as verbal or physical aggression toward people or objects.
 2. Reckless or self-destructive behavior.
 3. Hypervigilance.
 4. Exaggerated startle response.
 5. Problems with concentration.
 6. Sleep disturbance (e.g., difficulty falling or staying asleep or restless sleep).
- F. Duration of the disturbance (Criteria B, C, D, and E) is more than 1 month.

- G. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.
- H. The disturbance is not attributable to the physiological effects of a substance (e.g., medication, alcohol) or another medical condition.

Specify whether:

With dissociative symptoms: The individual's symptoms meet the criteria for posttraumatic stress disorder, and in addition, in response to the stressor, the individual experiences persistent or recurrent symptoms of either of the following:

1. **Depersonalization:** Persistent or recurrent experiences of feeling detached from, and as if one were an outside observer of, one's mental processes or body (e.g., feeling as though one were in a dream; feeling a sense of unreality of self or body or of time moving slowly).
2. **Derealization:** Persistent or recurrent experiences of unreality of surroundings (e.g., the world around the individual is experienced as unreal, dreamlike, distant, or distorted).

Note: To use this subtype, the dissociative symptoms must not be attributable to the physiological effects of a substance (e.g., blackouts, behavior during alcohol intoxication) or another medical condition (e.g., complex partial seizures).

Specify if:

With delayed expression: If the full diagnostic criteria are not met until at least 6 months after the event (although the onset and expression of some symptoms may be immediate).

Posttraumatic Stress Disorder for Children 6 Years and Younger

- A. In children 6 years and younger, exposure to actual or threatened death, serious injury, or sexual violence in one (or more) of the following ways:
 1. Directly experiencing the traumatic event(s).
 2. Witnessing, in person, the event(s) as it occurred to others, especially primary caregivers.

Note: Witnessing does not include events that are witnessed only in electronic media, television, movies, or pictures.

3. Learning that the traumatic event(s) occurred to a parent or caregiving figure.
- B. Presence of one (or more) of the following intrusion symptoms associated with the traumatic event(s), beginning after the traumatic event(s) occurred:
 1. Recurrent, involuntary, and intrusive distressing memories of the traumatic event(s).

Note: Spontaneous and intrusive memories may not necessarily appear distressing and may be expressed as play re-enactment.
 2. Recurrent distressing dreams in which the content and/or affect of the dream are related to the traumatic event(s).

Note: It may not be possible to ascertain that the frightening content is related to the traumatic event.
 3. Dissociative reactions (e.g., flashbacks) in which the child feels or acts as if the traumatic event(s) were recurring. (Such reactions may occur on a continuum, with the most extreme expression being a complete loss of awareness of present surroundings.) Such trauma-specific reenactment may occur in play.
 4. Intense or prolonged psychological distress at exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).
 5. Marked physiological reactions to reminders of the traumatic event(s).
 - C. One (or more) of the following symptoms, representing either persistent avoidance of stimuli associated with the traumatic event(s) or negative alterations in cognitions and mood associated with the traumatic event(s), must be present, beginning after the event(s) or worsening after the event(s):

Persistent Avoidance of Stimuli

1. Avoidance of or efforts to avoid activities, places, or physical reminders that arouse recollections of the traumatic event(s).

2. Avoidance of or efforts to avoid people, conversations, or interpersonal situations that arouse recollections of the traumatic event(s).

Negative Alterations in Cognitions

3. Substantially increased frequency of negative emotional states (e.g., fear, guilt, sadness, shame, confusion).
 4. Markedly diminished interest or participation in significant activities, including constriction of play.
 5. Socially withdrawn behavior.
 6. Persistent reduction in expression of positive emotions.
- D. Alterations in arousal and reactivity associated with the traumatic event(s), beginning or worsening after the traumatic event(s) occurred, as evidenced by two (or more) of the following:
1. Irritable behavior and angry outbursts (with little or no provocation) typically expressed as verbal or physical aggression toward people or objects (including extreme temper tantrums).
 2. Hypervigilance.
 3. Exaggerated startle response.
 4. Problems with concentration.
 5. Sleep disturbance (e.g., difficulty falling or staying asleep or restless sleep).
- E. The duration of the disturbance is more than 1 month.
- F. The disturbance causes clinically significant distress or impairment in relationships with parents, siblings, peers, or other caregivers or with school behavior.
- G. The disturbance is not attributable to the physiological effects of a substance (e.g., medication or alcohol) or another medical condition.

Specify whether:

With dissociative symptoms: The individual's symptoms meet the criteria for posttraumatic stress disorder, and the individual experiences persistent or recurrent symptoms of either of the following:

1. **Depersonalization:** Persistent or recurrent experiences of feeling detached from, and as if one were an outside ob-

server of, one's mental processes or body (e.g., feeling as though one were in a dream; feeling a sense of unreality of self or body or of time moving slowly).

2. **Derealization:** Persistent or recurrent experiences of unreality of surroundings (e.g., the world around the individual is experienced as unreal, dreamlike, distant, or distorted).

Note: To use this subtype, the dissociative symptoms must not be attributable to the physiological effects of a substance (e.g., blackouts) or another medical condition (e.g., complex partial seizures).

Specify if:

With delayed expression: If the full diagnostic criteria are not met until at least 6 months after the event (although the onset and expression of some symptoms may be immediate).

Acute Stress Disorder

308.3 (F43.0)

- A. Exposure to actual or threatened death, serious injury, or sexual violation in one (or more) of the following ways:
1. Directly experiencing the traumatic event(s).
 2. Witnessing, in person, the event(s) as it occurred to others.
 3. Learning that the event(s) occurred to a close family member or close friend. **Note:** In cases of actual or threatened death of a family member or friend, the event(s) must have been violent or accidental.
 4. Experiencing repeated or extreme exposure to aversive details of the traumatic event(s) (e.g., first responders collecting human remains, police officers repeatedly exposed to details of child abuse).
- Note:** This does not apply to exposure through electronic media, television, movies, or pictures, unless this exposure is work related.
- B. Presence of nine (or more) of the following symptoms from any of the five categories of intrusion, negative mood, dissociation,

avoidance, and arousal, beginning or worsening after the traumatic event(s) occurred:

Intrusion Symptoms

1. Recurrent, involuntary, and intrusive distressing memories of the traumatic event(s). **Note:** In children, repetitive play may occur in which themes or aspects of the traumatic event(s) are expressed.
2. Recurrent distressing dreams in which the content and/or affect of the dream are related to the event(s). **Note:** In children, there may be frightening dreams without recognizable content.
3. Dissociative reactions (e.g., flashbacks) in which the individual feels or acts as if the traumatic event(s) were recurring. (Such reactions may occur on a continuum, with the most extreme expression being a complete loss of awareness of present surroundings.) **Note:** In children, trauma-specific re-enactment may occur in play.
4. Intense or prolonged psychological distress or marked physiological reactions in response to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).

Negative Mood

5. Persistent inability to experience positive emotions (e.g., inability to experience happiness, satisfaction, or loving feelings).

Dissociative Symptoms

6. An altered sense of the reality of one's surroundings or oneself (e.g., seeing oneself from another's perspective, being in a daze, time slowing).
7. Inability to remember an important aspect of the traumatic event(s) (typically due to dissociative amnesia and not to other factors such as head injury, alcohol, or drugs).

Avoidance Symptoms

8. Efforts to avoid distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).
9. Efforts to avoid external reminders (people, places, conversations, activities, objects, situations) that arouse distressing

memories, thoughts, or feelings about or closely associated with the traumatic event(s).

Arousal Symptoms

10. Sleep disturbance (e.g., difficulty falling or staying asleep, restless sleep).
 11. Irritable behavior and angry outbursts (with little or no provocation), typically expressed as verbal or physical aggression toward people or objects.
 12. Hypervigilance.
 13. Problems with concentration.
 14. Exaggerated startle response.
- C. Duration of the disturbance (symptoms in Criterion B) is 3 days to 1 month after trauma exposure.

Note: Symptoms typically begin immediately after the trauma, but persistence for at least 3 days and up to a month is needed to meet disorder criteria.

- D. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.
- E. The disturbance is not attributable to the physiological effects of a substance (e.g., medication or alcohol) or another medical condition (e.g., mild traumatic brain injury) and is not better explained by brief psychotic disorder.

Adjustment Disorders

- A. The development of emotional or behavioral symptoms in response to an identifiable stressor(s) occurring within 3 months of the onset of the stressor(s).
- B. These symptoms or behaviors are clinically significant, as evidenced by one or both of the following:
1. Marked distress that is out of proportion to the severity or intensity of the stressor, taking into account the external context and the cultural factors that might influence symptom severity and presentation.

2. Significant impairment in social, occupational, or other important areas of functioning.
- C. The stress-related disturbance does not meet the criteria for another mental disorder and is not merely an exacerbation of a pre-existing mental disorder.
- D. The symptoms do not represent normal bereavement.
- E. Once the stressor or its consequences have terminated, the symptoms do not persist for more than an additional 6 months.

Specify whether:

309.0 (F43.21) With depressed mood: Low mood, tearfulness, or feelings of hopelessness are predominant.

309.24 (F43.22) With anxiety: Nervousness, worry, jitteriness, or separation anxiety is predominant.

309.28 (F43.23) With mixed anxiety and depressed mood: A combination of depression and anxiety is predominant.

309.3 (F43.24) With disturbance of conduct: Disturbance of conduct is predominant.

309.4 (F43.25) With mixed disturbance of emotions and conduct: Both emotional symptoms (e.g., depression, anxiety) and a disturbance of conduct are predominant.

309.9 (F43.20) Unspecified: For maladaptive reactions that are not classifiable as one of the specific subtypes of adjustment disorder.

Other Specified Trauma- and Stressor-Related Disorder

309.89 (F43.8)

This category applies to presentations in which symptoms characteristic of a trauma- and stressor-related disorder that cause clinically significant distress or impairment in social, occupational, or other important areas of functioning predominate but do not meet the full criteria for any of the disorders in the trauma- and stressor-related disorders diagnostic class. The other specified trauma- and stressor-related disorder category is used in situations in which the clinician

chooses to communicate the specific reason that the presentation does not meet the criteria for any specific trauma- and stressor-related disorder. This is done by recording "other specified trauma- and stressor-related disorder" followed by the specific reason (e.g., "persistent complex bereavement disorder").

Examples of presentations that can be specified using the "other specified" designation include the following:

1. **Adjustment-like disorders with delayed onset of symptoms that occur more than 3 months after the stressor.**
2. **Adjustment-like disorders with prolonged duration of more than 6 months without prolonged duration of stressor.**
3. **Ataque de nervios:** See "Glossary of Cultural Concepts of Distress" in the Appendix to DSM-5.
4. **Other cultural syndromes:** See "Glossary of Cultural Concepts of Distress" in the Appendix to DSM-5.
5. **Persistent complex bereavement disorder:** This disorder is characterized by severe and persistent grief and mourning reactions (see the chapter "Conditions for Further Study" in Section III of DSM-5).

Unspecified Trauma- and Stressor-Related Disorder

309.9 (F43.9)

This category applies to presentations in which symptoms characteristic of a trauma- and stressor-related disorder that cause clinically significant distress or impairment in social, occupational, or other important areas of functioning predominate but do not meet the full criteria for any of the disorders in the trauma- and stressor-related disorders diagnostic class. The unspecified trauma- and stressor-related disorder category is used in situations in which the clinician chooses *not* to specify the reason that the criteria are not met for a specific trauma- and stressor-related disorder, and includes presentations in which there is insufficient information to make a more specific diagnosis (e.g., in emergency room settings).